


# Lecture 8

## Some Discrete Probability Distributions II

### Lecture Information

Ceng272 *Statistical Computations* at April 12, 2010

Dr. Cem Özdoğan  
Computer Engineering Department  
Çankaya University




## 1 Negative Binomial and Geometric Distributions

Negative Binomial and  
Geometric  
Distributions

Poisson Distribution  
and the Poisson  
Process

## 2 Poisson Distribution and the Poisson Process



- **Negative binomial distribution (Geometric distribution):** the number of trial on which the first success occurs.
- **Poisson distribution:** the number of outcomes occurring during a given time interval or in a specified region.


Negative Binomial and  
Geometric  
Distributions

Poisson Distribution  
and the Poisson  
Process

## Negative Binomial and Geometric Distributions I

- Suppose, instead of performing a fixed or given number of trials, one performs independently a Bernoulli trial repeatedly until a desired number of successes is obtained, and then stop.
- Then, the question is that how many trials are required to get the desired number of successes?
- **Negative binomial experiments:** the  $k^{\text{th}}$  success occurs on the  $x^{\text{th}}$  trial.
- **Negative binomial random variable:** the number  $X$  of trials to produce  $k$  success in a negative binomial experiment.
- **Negative binomial distribution:** If repeated independent trials can result in a success with probability  $p$  and a failure with probability  $q = 1 - p$ , then the probability distribution of the random variable  $X$ , the number of the trial on which the  $k^{\text{th}}$  success occurs, is

$$b^*(x; k, p) = \binom{x-1}{k-1} p^k q^{x-k}, \quad x = k, k+1, k+2, \dots$$



## Negative Binomial and Geometric Distributions II

- **Example 5.17:** Suppose that team  $A$  has probability 0.55 of winning over the team  $B$  and both teams  $A$  and  $B$  face each other in an NBA 4-out-of-7 championship series.
  - i What is the probability that team  $A$  will win the series in six games?


$$b^*(x; k, p) = b^*(6; 4, 0.55) = \binom{5}{3} * 0.55^4 * (1 * 0.55)^{6-4} = 0.1853$$


- ii What is the probability that team  $A$  will win the series?

$$\begin{aligned} b^*(4; 4, 0.55) + b^*(5; 4, 0.55) + b^*(6; 4, 0.55) + b^*(7; 4, 0.55) \\ = 0.0915 + 0.1647 + 0.1853 + 0.1668 = 0.6083 \end{aligned}$$

- iii If both teams face each other in a regional play-off series and the winner is decided by winning three out of five games, what is the probability that team  $A$  will win a play-off?

$$\begin{aligned} b^*(3; 3, 0.55) + b^*(4; 3, 0.55) + b^*(5; 3, 0.55) \\ = 0.1664 + 0.2246 + 0.2021 = 0.5931 \end{aligned}$$






- **Why name negative binomial?** The binomial coefficient is defined even when  $n$  is negative (or is not an integer).

$$\binom{n}{k} = \frac{n(n-1)(n-2)\dots(n-k+1)}{k!}$$

$$(p+q)^n = \sum_{k=0}^{\infty} \binom{n}{k} p^k q^{n-k}$$


- Each term in the expansion of  $p^k(1-q)^{-k}$  corresponds to the value of  $b^*(x; k, p)$  for  $x = k, k+1, k+2, \dots$

$$1 = p^k * p^{-k} = p^k * (1-q)^{-k} = p^k * \sum_{x=0}^{\infty} \binom{-k}{x} (-q)^x$$



- Consider a binomial experiment to get a first success. This implies that for the case of that we encountered the number of failures prior to the first success.
- **Geometric Distribution:** If repeated independent trials can result in a success with probability  $p$  and a failure with probability  $q = 1 - p$ , then the probability distribution of the random variable  $X$ , the number of the trial on which the first success occurs, is

$$g(x; p) = b^*(x; 1, p) = pq^{x-1}, x = 1, 2, 3, \dots$$



- **Example:** Consider a problem of log in into a communication network. It is know that the probability of success rate,  $p = 0.35$  during busy hours.

- i Probability that one is able to log into the network at  $3^{rd}$  trial:

$$P(Y = 3) = (0.65)^2(0.35) = 0.1479.$$

- ii Probability that one is able to log into within 3 trials:

$$P(Y \leq 3) = P(Y = 1) + P(Y = 2) + P(Y = 3)$$

$$= 0.35 + 0.2275 + 0.1479 = 0.7254$$


- iii The average number of trials to get into the network:

$$E(Y) = 1/0.35 = 2.85.$$

That is, it takes about 3 times on average



## Negative Binomial and Geometric Distributions VII




- **Example 5.18:** In a certain manufacturing process it is known that, on the average, 1 in every 100 items is defective.
- What is the probability that the fifth item inspected is the first defective item found?

$$g(x; p) = g(5; 0.01) = 0.01 \times 0.99^4 = 0.0096$$

- **Example 5.19:** At “busy time” a telephone exchange is very near capacity, so callers have difficulty placing their calls.
- It may be of interest to know the number of attempts necessary in order to gain a connection.
- Suppose that let  $p = 0.05$  be the probability of a connection during busy time.
- We are interested in knowing the probability that 5 attempts are necessary for a successful call.

$$P(X = x) = g(5; 0.05) = 0.05 * 0.95^4 = 0.041$$




- **Theorem 5.4:**


The mean and variance of a random variable following the geometric distribution are

$$\mu = \frac{1}{p}, \quad \sigma^2 = \frac{1-p}{p^2}$$

- In the system of telephone exchange, trials occurring prior to a success represent a cost.
- A high probability of requiring a large of number of attempts is not beneficial to the scientists or engineers.



- **Poisson experiments:** Experiments yielding numerical values of a random variable  $X$ , the number of outcomes occurring during a given time interval or in a specified region.
- Examples:
  - the number of telephone calls per hour received by an office
  - the number of postponed baseball games due to rain
  - the number of field mice per acre
  - the number of typing error per page
- The examples of the random variables that are having Poisson probability distribution are usually rare events such as # of car accident, # of flood or hurricane occurrences.
- So the Poisson distribution provides the fundamental idea of the Law of Small Numbers (LSN).




Negative Binomial and  
Geometric  
Distributions

Poisson Distribution  
and the Poisson  
Process

- **Properties of Poisson Process:**
  - 1 The number of outcomes in one time interval or specified region is independent of the number that occurs in any other disjoint time interval or region of space. In this way we say that the Poisson process has no memory.
  - 2 The probability that a single outcome will occur during a very short time interval or in a small region is proportional to the length of the time interval or the size of the region and does not depend on the number of outcomes occurring outside this time interval or region.
  - 3 The probability that more than one outcome will occur in such a short time interval or fall in such a small region is negligible.

## Poisson Distribution and the Poisson Process III




- **Poisson Distribution:** The probability distribution of the Poisson random variable  $X$ , representing the number of outcomes occurring in a given time interval or specified region denoted by  $t$ , is (mean number  $\mu = \lambda t$ )

$$p(x; \lambda t) = \frac{e^{-\lambda t} (\lambda t)^x}{x!}$$

- where  $\lambda$  is the average number of outcomes per unit time or region, and  $e = 2.71828$ .
- **Example 5.20:** During a laboratory experiment the average number of radioactive particles passing through a counter in 1 millisecond is 4.
- What is the probability that 6 particles enter the counter in a given millisecond? (see Table A.2)

$$\begin{aligned} p(6; 4) &= \frac{e^{-4}(4)^6}{6!} = \sum_{x=0}^6 p(x; 4) - \sum_{x=0}^5 p(x; 4) \\ &= 0.8893 - 0.7851 = 0.1042 \end{aligned}$$

# Poisson Distribution and the Poisson Process IV



**TABLE A.2** Poisson Probability Sums  $\sum_{x=0}^r p(x; \mu)$


| $r$ | $\mu$  |        |        |        |        |        |        |        |        |
|-----|--------|--------|--------|--------|--------|--------|--------|--------|--------|
|     | 0.1    | 0.2    | 0.3    | 0.4    | 0.5    | 0.6    | 0.7    | 0.8    | 0.9    |
| 0   | 0.9048 | 0.8187 | 0.7408 | 0.6730 | 0.6065 | 0.5488 | 0.4966 | 0.4493 | 0.4066 |
| 1   | 0.9953 | 0.9825 | 0.9631 | 0.9384 | 0.9098 | 0.8781 | 0.8442 | 0.8088 | 0.7725 |
| 2   | 0.9998 | 0.9989 | 0.9964 | 0.9921 | 0.9856 | 0.9769 | 0.9659 | 0.9526 | 0.9371 |
| 3   | 1.0000 | 0.9999 | 0.9997 | 0.9992 | 0.9982 | 0.9966 | 0.9942 | 0.9909 | 0.9865 |
| 4   |        | 1.0000 | 1.0000 | 0.9999 | 0.9998 | 0.9996 | 0.9992 | 0.9986 | 0.9977 |
| 5   |        |        |        | 1.0000 | 1.0000 | 1.0000 | 0.9999 | 0.9998 | 0.9997 |
| 6   |        |        |        |        |        |        | 1.0000 | 1.0000 | 1.0000 |
| ∞   |        |        |        |        |        |        |        |        |        |

| $r$ | $\mu$  |        |        |        |        |        |        |        |        |  |
|-----|--------|--------|--------|--------|--------|--------|--------|--------|--------|--|
|     | 1.0    | 1.5    | 2.0    | 2.5    | 3.0    | 3.5    | 4.0    | 4.5    | 5.0    |  |
| 0   | 0.3679 | 0.2231 | 0.1353 | 0.0821 | 0.0498 | 0.0302 | 0.0183 | 0.0111 | 0.0067 |  |
| 1   | 0.7358 | 0.5578 | 0.4060 | 0.2873 | 0.1991 | 0.1359 | 0.0916 | 0.0611 | 0.0404 |  |
| 2   | 0.9197 | 0.8088 | 0.6767 | 0.5438 | 0.4232 | 0.3208 | 0.2381 | 0.1736 | 0.1247 |  |
| 3   | 0.9810 | 0.9344 | 0.8571 | 0.7576 | 0.6472 | 0.5366 | 0.4335 | 0.3423 | 0.2650 |  |
| 4   | 0.9963 | 0.9814 | 0.9473 | 0.8912 | 0.8153 | 0.7254 | 0.6288 | 0.5321 | 0.4405 |  |
| 5   | 0.9994 | 0.9955 | 0.9834 | 0.9580 | 0.9161 | 0.8576 | 0.7851 | 0.7029 | 0.6160 |  |
| 6   | 0.9999 | 0.9991 | 0.9955 | 0.9858 | 0.9665 | 0.9347 | 0.8893 | 0.8311 | 0.7622 |  |
| 7   | 1.0000 | 0.9998 | 0.9989 | 0.9958 | 0.9881 | 0.9733 | 0.9489 | 0.9134 | 0.8666 |  |
| 8   |        | 1.0000 | 0.9998 | 0.9989 | 0.9962 | 0.9901 | 0.9786 | 0.9597 | 0.9319 |  |
| 9   |        |        | 1.0000 | 0.9997 | 0.9989 | 0.9967 | 0.9919 | 0.9829 | 0.9682 |  |
| 10  |        |        |        | 0.9999 | 0.9997 | 0.9990 | 0.9972 | 0.9933 | 0.9863 |  |
| 11  |        |        |        |        | 1.0000 | 0.9999 | 0.9997 | 0.9976 | 0.9945 |  |
| 12  |        |        |        |        |        | 1.0000 | 0.9999 | 0.9992 | 0.9980 |  |
| 13  |        |        |        |        |        |        | 1.0000 | 0.9997 | 0.9993 |  |
| 14  |        |        |        |        |        |        |        | 1.0000 | 0.9998 |  |
| 15  |        |        |        |        |        |        |        |        | 1.0000 |  |
| 16  |        |        |        |        |        |        |        |        |        |  |

**Figure:** Poisson Probability Sums  $P(x; \mu) = \sum_{x=0}^r p(x; \mu)$ .

## Poisson Distribution and the Poisson Process V




- **Example 5.21:** Ten is the average number of oil tankers arriving each day at a certain port city.
- The facilities at the port can handle at most 15 tankers per day.
- What is the probability that on a given day tankers have to be turned away?

$$\begin{aligned}P(X > 15) &= 1 - P(X \leq 15) = 1 - \sum_{x=0}^{15} p(x; 10) \\ &= 1 - 0.9513 = 0.0487\end{aligned}$$

- **Theorem 5.5:**

The mean and variance of the Poisson distribution  $p(x; \lambda t)$  both have the value  $\lambda t$ . (Proof is in Appendix A.26)

- **Example:** In Example 5.20,  
 $\lambda t = 4 \Rightarrow \mu = \sigma^2 = 4, \sigma = 2 \Rightarrow \mu \pm 2\sigma = 4 \pm 2 * 2$ .
- Using Chebyshev's theorem, we conclude that at least 3/4 of the time the number of radioactive particles entering the counter will be anywhere from 0 to 8 during a given millisecond.



- **The Poisson Distribution As a Limiting Form of the Binomial: Theorem 5.6:**

Let  $X$  be a binomial random variable with probability distribution  $b(x; n, p)$ . When  $n \rightarrow \infty$ ,  $p \rightarrow 0$ , and  $\mu = np$  remains constant,



$$b(x; n, p) \xrightarrow{n \rightarrow \infty} p(x; \mu)$$

(Proof is in Appendix A.27)

- If  $p \rightarrow 1$ , we can change  $p$  to a value close to 0 by interchanging what we have defined to be a success and a failure.




## Poisson Distribution and the Poisson Process VII



**Figure:** Poisson density functions for different means.

The shape becomes more symmetric as the mean grows large.




- **Example 5.22:** The probability of an accident in a certain industrial facility on any given day is 0.005 and accidents are independent of each other.
  - i What is the probability that in any given period of 400 days there will be an accident on one day?

$$n = 400, p = 0.005 \Rightarrow \mu = np = 2$$

$$P(X = 1) = e^{-2}2^1 = 0.271$$

- ii What is the probability that there are at most three days with an accident?

$$P(X \leq 3) = \sum_{x=0}^3 \frac{e^{-2}2^x}{x!} = 0.857$$



- **Example 5.23:** In a manufacturing process where glass products are produced, defects or bubbles occur, occasionally rendering the piece undesirable for marketing.
- It is known that, on average, 1 in every 1000 of these items produced has one or more bubbles.
- What is the probability that a random sample of 8000 will yield fewer than 7 items processing bubbles?
- Solution:

$$n = 8000, p = 0.001 \Rightarrow \mu = np = 8$$

$$P(X \leq 7) = \sum_{x=0}^6 b(x; 8000, 0.001) \approx \sum_{x=0}^6 p(x; 8) = 0.3134$$